

## Comparison of frequency-selective properties of meteor burst channel at the beginning and at the end of radio reflections from meteor trails

Sulimov A., Safiullina A., Karpov A.

Kazan Federal University, 420008, Kremlevskaya 18, Kazan, Russia

---

### Abstract

© 2017 IEEE. Meteor burst propagation is based on scattering of radio waves from the ionized trails left by fast meteor particles. One of the promising applications of meteor radio propagation is development of Meteor Synchronization Systems (MSSs) for synchronizing two remote stations with potentially subnanosecond accuracy. Frequency-selective properties of meteor burst channel limit potential accuracy of coherent multi-carrier MSS systems. To adapt to frequency distortions of timing signals, a frequency response of the channel should be appropriately considered. Continuous expansion of the meteor trail plasma due to ambipolar diffusion provokes significant changes in frequency-selective properties of the channel at the end of detection of radio reflections. This should be taken into account to assess a minimum channel bandwidth correctly. On the basis of rigorous solution to the problem of diffraction of radio waves on a meteor trail, we perform a numerical simulation of the amplitude-frequency and phase-frequency responses of meteor burst channel. A comparison of the frequency-selective properties of the channel at the beginning and at the end of the signal detection is presented for the cases of radio reflections from underdense and overdense meteor trails. The frequency responses obtained with the diffraction approach are compared with the results of the classical (approximate) theory of a radio reflection from meteor trail.

<http://dx.doi.org/10.1109/RSEMW.2017.8103589>

---

### Keywords

Diffraction of radio waves, Frequency response of channel, Meteor burst propagation, Meteor radio reflection

### References

- [1] J.D. Oetting, "An analysis of meteor burst communications for military applications, " IEEE Trans. on comm, vol. 28, no. 9, pp. 1591- 1601, 1980.
- [2] V.A. Korneyev, Time & Frequency coordination using unsteady, variable-precision measurements in meteor-burst channel, Proc. 2003 IEEE Int. Freq. Cont. Symp. and PDA Exhibition Jointly with the 17th EFTF, pp. 285-289, Tampa (USA), 4-8 May 2003.
- [3] L.A. Epictetov, "Application of Meteor Burst Equipment for High Precision Comparisons of Time and Frequency Standards, " Proc.of 7th European Frequency and Time Forum (EFTF'), pp.413-416, Neuchatel (Switzerland), 16-18 March 1993.

- [4] T.R. Kaiser, "Theory of radio reflections from meteor trails: I, " Philosophical Magazine: Series 7, vol. 43, pp.1-32, 1952.
- [5] P.S. Cannon, "Polarization rotation in meteor-burst communication systems, " Radio Science, vol. 21, no. 3, pp. 501-510, May-June 1986.
- [6] K. C. Yeh, "Note on the geometry of the earth magnetic field useful to Faraday effect experiments, " J. of Geophys. Res., vol.65, no.10, p.3209, 1960.
- [7] R.G. Khuziyashev, "Calculation of the amplitude and phase characteristics of a signal scattered obliquely off a meteor trail, " Radiophysics and Quantum Electronics, vol. 27, 9, pp. 778-782, 1984.
- [8] D. W. R. McKinley, "Meteor science and engineering, " McGraw-Hill, 1961.
- [9] A. Karpov, "The computer model KAMET: The new generation version, " Proc. Meteoroids 2001 Conf., pp.367-370, Kiruna (Sweden), 6- 10 Aug, 2001.
- [10] R.I. Jr. Desourdis, "Nonreciprocity of meteor scatter radio links, " Proc. Ionospheric Effects Symposium (IES), pp.165-173, 4-6 May 1993.
- [11] J.A. Weitzen, "Characterizing the Multipath and Doppler Spreads of the High-Latitude Meteor-burst Communication Channel, " IEEE Trans. on Comm., vol. 35, pp.1050-1058, 1987.
- [12] K.J. Ellis, "Impulse response measurements on a 500-km meteor burst link, " Radio Science, vol. 34, no. 6, pp. 1525-1536, 1999.